

321-POLYGON-AVOIDING PERMUTATIONS AND CHEBYSHEV POLYNOMIALS

TOUFIK MANSOUR

LaBri, Université Bordeaux I,
351 cours de la Libération, 33405 Talence Cedex
toufik@labri.fr

ZVEZDELINA STANKOVA

Mills College, Oakland, CA
stankova@mills.edu

ABSTRACT. A 321 - k -gon-avoiding permutation π avoids 321 and the following four patterns:

$$\begin{aligned} &k(k+2)(k+3) \cdots (2k-1)1(2k)23 \cdots (k+1), \quad k(k+2)(k+3) \cdots (2k-1)(2k)123 \cdots (k+1), \\ &(k+1)(k+2)(k+3) \cdots (2k-1)1(2k)23 \cdots k, \quad (k+1)(k+2)(k+3) \cdots (2k-1)(2k)123 \cdots k. \end{aligned}$$

The 321 - k -gon-avoiding permutations were introduced and studied by Billey and Warrington [BW] as a class of elements of the symmetric group whose Kazhdan-Lusztig, Poincaré polynomials, and the singular loci of whose Schubert varieties have fairly simple formulas and descriptions. Stankova and West [SW] gave an exact enumeration in terms of linear recurrences with constant coefficients for the cases $k = 2, 3, 4$. In this paper, we extend these results by finding an explicit expression for the generating function for the number of 321 - k -gon-avoiding permutations on n letters. The generating function is expressed via Chebyshev polynomials of the second kind.

1. INTRODUCTION

Definition 1. Let $\alpha \in S_n$ and $\tau \in S_k$ be two permutations. Then α *contains* τ if there exists a subsequence $1 \leq i_1 < i_2 < \cdots < i_k \leq n$ such that $(\alpha_{i_1}, \dots, \alpha_{i_k})$ is order-isomorphic to τ ; in such a context τ is usually called a *pattern*; α *avoids* τ , or is τ -*avoiding*, if α does not contain such a subsequence. The set of all τ -avoiding permutations in S_n is denoted by $S_n(\tau)$. For a collection of patterns T , α *avoids* T if α avoids all $\tau \in T$; the corresponding subset of S_n is denoted by $S_n(T)$.

While the case of permutations avoiding a single pattern has attracted much attention, the case of multiple pattern avoidance remains less investigated. In particular, it is natural to consider permutations avoiding pairs of patterns τ_1, τ_2 . This problem was solved completely for $\tau_1, \tau_2 \in S_3$ (see [SS]), for $\tau_1 \in S_3$ and $\tau_2 \in S_4$ (see [W]), and for $\tau_1, \tau_2 \in S_4$ (see [Bo1, Km] and references therein). Several recent papers [CW, MV1, Kr, MV2, MV3, MV4] deal with the case $\tau_1 \in S_3, \tau_2 \in S_k$ for various pairs τ_1, τ_2 . The tools involved in these papers include Fibonacci numbers, Catalan numbers, Chebyshev polynomials, continued fractions, and Dyck words, e.g. in [MV2]:

Theorem 1.1 (Mansour, Vainshtein). *Let $U_m(\cos \theta) = \sin(m+1)\theta / \sin \theta$ be the Chebyshev polynomial of the second kind. When $2 \leq d+1 \leq k$, the generating function for the number of permutations in*

$S_n(321, (d+1) \cdots k12 \cdots d)$ is given by

$$\frac{U_{k-1}\left(\frac{1}{2\sqrt{x}}\right)}{\sqrt{x}U_k\left(\frac{1}{2\sqrt{x}}\right)}.$$

Recently, a special class of restricted permutations has arisen in representation theory.

Definition 2. A permutation π is k -gon-avoiding if it avoids each pattern in the set $\mathfrak{P}_k$:

$$\left\{ \begin{aligned} &k(k+2)(k+3) \cdots (2k-1)1(2k)23 \cdots (k+1), & k(k+2)(k+3) \cdots (2k-1)(2k)123 \cdots (k+1), \\ &(k+1)(k+2)(k+3) \cdots (2k-1)1(2k)23 \cdots k, & (k+1)(k+2)(k+3) \cdots (2k-1)(2k)123 \cdots k \end{aligned} \right\}.$$

We say that π is a 321- k -gon-avoiding permutation if it is both k -gon-avoiding and 321-avoiding. The number of 321- k -gon-avoiding permutations in S_n is denoted by $f_k(n)$. The corresponding generating function is $f_k(x) = \sum_{n \geq 0} f_k(n)x^n$.




FIGURE 1. $\mathfrak{B}_5$: all four 5-gons

Note that $f_k(n) = \frac{1}{n+1} \binom{2n}{n}$ for $n \in [0, 2k-1]$, as these count the permutations in $S_n(321)$ (see [Kn]).

Billey and Warrington [BW] introduced the 321-4-gon-avoiding (or 321-hexagon-avoiding) permutations as a class in S_n whose Kazhdan-Lusztig and Poincaré polynomials, and the singular loci of whose Schubert varieties have fairly simple formulas and descriptions. Upon their request, Stankova and West [SW] presented an exact enumeration for the cases $k = 2, 3, 4$ by using generating trees, the symmetries in the set of the $\mathfrak{P}_k$, and the structure of the 321-avoiding permutations via Schensted's 321-subsequences decomposition.

Theorem 1.2 (Stankova, West). *For $k = 2, 3, 4$, the sequences $f_k(n)$ satisfy the recursive relations*

$$\begin{aligned} f_4(n) &= 6f_4(n-1) - 11f_4(n-2) + 9f_4(n-3) - 4f_4(n-4) - 4f_4(n-5) + f_4(n-6), \quad n \geq 6; \\ f_3(n) &= 4f_3(n-1) - 4f_3(n-2) + 3f_3(n-3) + f_3(n-4) - f_3(n-5), \quad n \geq 5; \\ f_2(n) &= 3f_2(n-1) - 3f_2(n-2) + f_2(n-3) = (n-1)^2 + 1, \quad n \geq 3. \end{aligned}$$

In this paper we present an approach to the study of 321- k -gon-avoiding permutations in S_n which generalizes the methods in [SW] and [MV3]. As a consequence, we extend the results in [SW] to all 321- k -gon-avoiding permutations, and derive a number of other related results. The main theorem of the paper is formulated as follows.

Theorem 1.3. For $k \geq 3$ and $s \geq 1$, define $L_n^k(s) = \sum_{j=0}^s (-1)^j \binom{s-j}{j} f_k(n-j)$. When $n \geq 2k$, this sequence satisfies the linear recursive relation with constant coefficients:

$$L_n^k(2k-2) = L_{n-1}^k(2k-2) + L_{n-3}^k(2k-5) + L_{n-3}^k(2k-4) + L_{n-4}^k(2k-5) + L_{n-4}^k(2k-4).$$

Corollary 1.4. For $k \geq 3$,

$$f_k(x) = \frac{(1+x^2-x^4)U_{2k-3}\left(\frac{1}{2\sqrt{x}}\right) - \sqrt{x}(1+x^3)U_{2k-4}\left(\frac{1}{2\sqrt{x}}\right)}{\sqrt{x}\left[(1+x^2-x^4)U_{2k-2}\left(\frac{1}{2\sqrt{x}}\right) - \sqrt{x}(1+x^3)U_{2k-3}\left(\frac{1}{2\sqrt{x}}\right)\right]}.$$

The proofs of Theorem 1.3 and Corollary 1.4 are presented in Section 2. Note that Corollary 1.4 implies the previously known results for the cases $k = 3, 4$ (see [SW]):

$$\begin{aligned} f_3(x) &= \frac{1-3x+2x^2-2x^3-2x^4+3x^5}{1-4x+4x^2-3x^3-x^4+5x^5-x^6}, \\ f_4(x) &= \frac{1-5x+7x^2-5x^3+x^4+7x^5-4x^6}{1-6x+11x^2-9x^3+4x^4+8x^5-9x^6+x^7}. \end{aligned}$$

In Section 3, we describe several generalizations of Theorem 1.3 and Corollary 1.4, following similar arguments from their proofs.

2. PROOF THEOREM 1.3

2.1. Refinement of the numbers $f_k(n)$.

Definition 3. For m, n with $1 \leq m \leq n$ and distinct $i_1, i_2, \dots, i_m \in \mathbb{N}$, we denote by $f_k(n; i_1, \dots, i_m)$ the number of 321- k -gon-avoiding permutations $\pi \in S_n$ which start with $i_1 i_2 \dots i_m$: $\pi_1 \pi_2 \dots \pi_m = i_1 i_2 \dots i_m$. The corresponding subset of S_n is denoted by $\mathcal{F}_k(n; i_1, \dots, i_m)$.

Here follow basic properties of the numbers $f_k(n; i_1, \dots, i_m)$, easily deduced from the definitions.

Lemma 2.1. Let $n \geq 3$, $1 \leq m \leq n$ and $1 \leq i_1 < i_2 < \dots < i_m \leq n$.

(a) If $m \leq n-2$ and $3 \leq i_1$, then $f_k(n; i_1, \dots, i_m, j) = 0$ for $2 \leq j \leq i_m-1$. Consequently,

$$f_k(n; i_1, \dots, i_m) = f_k(n; i_1, \dots, i_m, 1) + \sum_{j=i_m+1}^n f_k(n; i_1, \dots, i_m, j).$$

(b) If $m \leq k-3$ and $2 \leq i_1$, then $f_k(n; i_1, \dots, i_m, 1) = f_k(n-1; i_1-1, \dots, i_m-1)$.

(c) If $i_1 \leq k-1$, then $f_k(n; i_1, \dots, i_m) = f_k(n-1; i_2-1, \dots, i_m-1)$.

Proof. For (a), observe that if $\pi \in \mathcal{F}_k(n; i_1, \dots, i_m, j)$ then the entries $i_m, j, 1$ give an occurrence of 321 in π . For the second part of (a), consider the entries π_{m+1} of π . Again, avoiding 321 forces $\pi_{m+1} = 1$ or $\pi_{m+1} > i_m$.

For (b), denote by π' the permutation obtained from π by deleting its smallest entry and decreasing all other entries by 1. Then $\pi \in \mathcal{F}_k(n; i_1, \dots, i_m, 1)$ if and only if $\pi' \in \mathcal{F}_k(n-1; i_1, \dots, i_m)$, since entry 1 placed as in (b) cannot be used in an occurrence of 321 or $\tau \in \mathfrak{P}_k$ in π .

For (c), observe that if $\pi_1\pi_2\cdots\pi_m = i_1i_2\cdots i_m$ then the entry i_1 cannot appear in any occurrences of $\tau \in \mathfrak{P}_k$; further, if there is an occurrence xyz of 321 such that $x = i_1$ then there is an occurrence i_2yx of 321 in π . $\square$

Lemma 2.1 implies an explicit formula for $f_k(n; s)$ for the first values of s .

Proposition 2.2. *For $1 \leq s \leq \min\{k-1, n\}$,*

$$f_k(n; s) = \sum_{j=0}^{s-1} (-1)^j \binom{s-1-j}{j} f_k(n-1-j).$$

Proof. By Lemma 2.1(c)-(a) the proposition holds for $s = 1, 2$. For $s \geq 3$, Lemma 2.1(b) says

$$\begin{aligned} f_k(n; s) &= f_k(n; s, 1) + \sum_{j=s+1}^n f_k(n; s, j) \\ \Rightarrow f_k(n; s) &= f_k(n-1; s-1) + \sum_{j=s+1}^n f_k(n-1; j-1) = \sum_{j=s-1}^{n-1} f_k(n-1; j). \end{aligned}$$

Equivalently,

$$(2.1) \quad f_k(n; s) = f_k(n-1) - \sum_{j=1}^{s-2} f_k(n-1; j).$$

Using induction on s , we assume that the proposition holds for all $1 \leq j \leq s-1$. Then (2.1) yields

$$f_k(n; s) = f_k(n-1) - \sum_{j=1}^{s-2} \sum_{i=0}^{j-1} (-1)^i \binom{j-1-i}{i} f_k(n-2-i).$$

From the familiar equality $\binom{1}{a} + \binom{2}{a} + \cdots + \binom{b}{a} = \binom{b+1}{a+1}$ we obtain for all $2 \leq s \leq k-1$

$$\begin{aligned} f_k(n; s) &= f_k(n-1) + \sum_{j=1}^{s-1} (-1)^j \binom{s-1-j}{j} f_k(n-1-j) \\ &= \sum_{j=0}^{s-1} (-1)^j \binom{s-1-j}{j} f_k(n-1-j). \quad \square \end{aligned}$$

Next we introduce objects $A_d(n, m)$ which organize suitably the information about the numbers $f_k(n; i_1, i_2, \dots, i_m)$ and play an important role in the proof of the main result.

Definition 4. For $1 \leq d \leq n+1-m$ and $1 \leq m \leq n$ set

$$A_d(n, m) = \sum_{d \leq i_1 < i_2 < \cdots < i_m \leq n} f_k(n; i_1, \dots, i_m).$$

In the following subsections 2.2–2.3 we derive two expressions for $A_k(n, k-1)$, compare them in subsection 2.4, and thus complete the proof of Theorem 1.3.

2.2. First expression for $A_k(n, k-1)$. The numbers $A_d(n, m)$ satisfy the following recurrence.

Lemma 2.3. For $2 \leq d \leq k$ and $1 \leq m \leq \min\{k-1, n\}$,

$$A_d(n, m) = A_{d-1}(n, m) - A_{d-1}(n-1, m-1).$$

PROOF. By (2.1), for all $2 \leq d \leq k$ we have:

$$A_d(n, m) = A_{d-1}(n, m) - \sum_{d-1=i_2 < \dots < i_m \leq n} f_k(n; d-1, i_2, \dots, i_m).$$

Lemma 2.1(c) and Definition 4 imply

$$\begin{aligned} A_d(n, m) &= A_{d-1}(n, m) - \sum_{d-1=i_2 < i_3 < \dots < i_m \leq n-1} f_k(n-1; i_2, i_3, \dots, i_m) \\ &= A_{d-1}(n, m) - A_{d-1}(n-1, m-1). \quad \square \end{aligned}$$

We next find an explicit expression for $A_1(n, m)$ in terms of $f_k(n)$.

Lemma 2.4. Let $1 \leq m \leq \min\{k-1, n\}$. Then

$$A_1(n, m) = \sum_{j=0}^m (-1)^j \binom{m-j}{j} f_k(n-j).$$

Proof. For $m = 1$, $A_1(n, 1) = \sum_{1 \leq i_1 \leq n} f_k(n; i_1) = f_k(n)$, which equals the required expression. Assume the lemma for m and all appropriate n . Comparing the $(m+1)^{\text{st}}$ entry j of π with i_m ,

$$A_1(n, m) = A_1(n, m+1) + \sum_{1 \leq i_1 < i_2 < \dots < i_m \leq n} \sum_{j=1}^{i_m-1} f_k(n; i_1, i_2, \dots, i_m, j).$$

For the summation part on the right-hand side, avoidance of (321) implies that all numbers $1, 2, \dots, j-1$ appear before the entry j , and hence $j \leq m$. From Lemma 2.1,

$$\begin{aligned} A_1(n, m) &= A_1(n, m+1) + \sum_{j=1}^m \sum_{j+1 \leq i_j < i_{j+1} < \dots < i_m \leq n} f_k(n; 1, 2, \dots, j-1, i_j, i_{j+1}, \dots, i_m, j) \\ &= A_1(n, m+1) + \sum_{j=1}^m \sum_{2 \leq i_j < i_{j+1} < \dots < i_m \leq n-j+1} f_k(n-j+1; i_j, i_{j+1}, \dots, i_m, 1) \\ &= A_1(n, m+1) + \sum_{j=1}^m \sum_{1 \leq i_j < i_{j+1} < \dots < i_m \leq n-j} f_k(n-j; i_j, i_{j+1}, \dots, i_m) \\ &= A_1(n, m+1) + \sum_{j=1}^m A_1(n-j, m-j+1) = \sum_{j=0}^m A_1(n-j, m-j+1) \end{aligned}$$

Applying the above to $A_1(n-1, m-1)$, subtracting the results, and using the induction hypothesis, we arrive at

$$\begin{aligned}
A_1(n, m) &= A_1(n, m+1) + A_1(n-1, m-1) \\
\Rightarrow A_1(n, m+1) &= A_1(n, m) - A_1(n-1, m-1) \\
&= \sum_{j=0}^m (-1)^j \binom{m-j}{j} f_k(n-j) - \sum_{j=0}^{m-1} (-1)^j \binom{m-j-1}{j} f_k(n-1-j) \\
&= \sum_{j=0}^{m+1} (-1)^j \binom{m-j+1}{j} f_k(n-j). \quad \square
\end{aligned}$$

Lemmas 2.3–2.4 can be combined to derive the following explicit expression for $A_d(n, m)$, which is easily proven by induction on d .

Corollary 2.5. *Let $1 \leq d \leq k$, $1 \leq m \leq \min\{k-1, n\}$. Then*

$$A_d(n, m) = \sum_{j=0}^{d+m-1} (-1)^j \binom{d+m-1-j}{j} f_k(n-j) = L_n^k(d+m-1).$$

In particular,

$$A_k(n, k-1) = \sum_{j=0}^{2k-2} (-1)^j \binom{2k-2-j}{j} f_k(n-j) = L_n^k(2k-2).$$

2.3. Second expression for $A_k(n, k-1)$. We start by introducing three objects related to $A_d(n, m)$.

Definition 5. For $1 \leq d \leq n-m+1$ and $1 \leq m \leq n$ set

$$\begin{aligned}
B_d(n, m) &= \sum_{d+1 \leq i_1 < i_2 < \dots < i_m \leq n} f_k(n; d, i_1, \dots, i_m); \\
C_d(n, m) &= \sum_{d \leq i_1 < i_2 < \dots < i_m \leq n} f_k(n; i_1, \dots, i_m, 1); \\
D_d(n, m) &= \sum_{d+1 \leq i_1 < i_2 < \dots < i_m \leq n} f_k(n; d, i_1, \dots, i_m, 1).
\end{aligned}$$

Note that by Lemma 2.1(a), for $k \geq 2$:

$$(2.2) \quad A_k(n, k-1) = A_k(n, k) + C_k(n, k-1).$$

The following Propositions 2.6–2.7 describe $A_k(n, k)$ and $C_k(n, k-1)$ in terms of $f_k(n)$.

Proposition 2.6. *Let $n \geq k-1$. Then*

- (a) $C_k(n, k-1) = C_{k-1}(n-1, k-1) + A_{k-1}(n-3, k-2);$
- (b) $C_k(n, k-1) = C_k(n-1, k-1) + A_{k-2}(n-3, k-2) + A_{k-1}(n-3, k-2).$

Proof. For (a), similarly to Lemma 2.1(b) (with $k \geq 3$), we have

$$(2.3) \quad C_k(n, k-1) = \sum_{k \leq i_1 < \dots < i_{k-1} \leq n} f_k(n; i_1, \dots, i_{k-1}, 1, 2) + \sum_{k \leq i_1 < \dots < i_{k-1} < i_k \leq n} f_k(n; i_1, \dots, i_{k-1}, 1, i_k).$$

If π starts with $i_1, \dots, i_{k-1}, 1, 2$ as in the first sum in (2.3), then the entry 2 cannot participate in an occurrence of 321 or of $\tau \in \mathfrak{P}_k$. Hence

$$(2.4) \quad f_k(n; i_1, \dots, i_{k-1}, 1, 2) = f_k(n-1; i_1-1, \dots, i_{k-1}-1, 1).$$

For the second sum in (2.3), if π starts with $i_1, \dots, i_{k-1}, 1, i_k$, avoidance of 321 and both

$$(k+1)(k+2)(k+3) \cdots (2k-1)1(2k)23 \cdots k \text{ and } k(k+2)(k+3) \cdots (2k-1)1(2k)23 \cdots (k+1)$$

implies $i_1 = k$ and $i_2 = k+1$. Now, if π starts with $k, k+1, i_3, \dots, i_{k-1}, 1, i_k$ where $k+2 \leq i_3 < \dots < i_k \leq n$, then note that no occurrence of $\tau \in \mathfrak{P}_k$ can contain the entries k or $k+1$; further, an occurrence of 321 containing k will exist in π if and only if there is such an occurrence containing $k+1$. Using this and Lemma 2.1,

$$\begin{aligned} f_k(n; k, k+1, i_3, \dots, i_{k-1}, 1, i_k) &= f_k(n-1; k, i_3-1, \dots, i_{k-1}-1, 1, i_k-1) \\ &= f_k(n-2; k-1, i_3-2, \dots, i_{k-1}-2, i_k-2) \\ &= f_k(n-3; i_3-3, \dots, i_{k-1}-3, i_k-3). \end{aligned}$$

Combining the last equality with (2.3), (2.4) and the definitions of $C_d(n, m)$ and $A_d(n, m)$, we obtain the desired equality

$$C_k(n, k-1) = C_{k-1}(n-1, k-1) + A_{k-1}(n-3, k-2).$$

For (b), by definitions of $C_d(n, m)$ and $D_d(n, m)$, we have

$$C_{k-1}(n-1, k-1) = C_k(n-1, k-1) + D_{k-1}(n-1, k-2).$$

Combining with (a), it is enough to show $D_{k-1}(n-1, k-2) = A_{k-2}(n-3, k-2)$. To this end, note that if $\pi \in \mathcal{F}_k(n-1; k-1, i_1, \dots, i_{k-2}, 1)$ where $k+1 \leq i_1 < \dots < i_{k-2} \leq n-1$, then by Lemma 2.1

$$f_k(n-1; k-1, i_1, \dots, i_{k-2}, 1) = f_k(n-3; i_1-2, \dots, i_{k-2}-2).$$

By definitions of $D_d(n, m)$ and $A_d(n, m)$, we obtain the required equality. $\square$

Proposition 2.7. *Let $n \geq k-1$. The sequences A_k, B_k, C_k and D_k satisfy the relations:*

- (a) $A_k(n, k) = B_k(n-1, k-2)$;
- (b) $B_k(n-1, k-2) = D_k(n-1, k-2) + B_k(n-1, k-1)$;
- (c) $B_k(n-1, k-1) = B_k(n-2, k-2)$;
- (d) $D_k(n-1, k-2) = A_{k-2}(n-4, k-2) + A_{k-1}(n-4, k-2)$;
- (e) $A_k(n, k) - A_k(n-1, k) = A_{k-2}(n-4, k-2) + A_{k-1}(n-4, k-2)$.

Proof. For (a), if $\pi \in F_k(n; i_1, i_2, \dots, i_k)$ so that $k \leq i_1 < i_2 < \dots < i_k \leq n$, then avoidance of 321,

$$(k+1)(k+2)(k+3) \cdots (2k-1)(2k)123 \cdots k \text{ and } k(k+2)(k+3) \cdots (2k-1)(2k)123 \cdots (k+1)$$

implies $i_1 = k$ and $i_2 = k+1$. Since no occurrence of $\tau \in \mathfrak{P}_k$ in π can contain both entries k and $k+1$, it follows that $f_k(n; k, k+1, i_3, \dots, i_k) = f_k(n-1; k, i_3-1, \dots, i_k-1)$, and hence (a).

Part (b) is obtained similarly as in Lemma 2.1(b).

For (c), if $\pi \in F_k(n-1; k, i_1, \dots, i_{k-1})$ where $k+1 \leq i_1 < \dots < i_{k-1} \leq n-1$, then avoidance of

$$321 \text{ and } k(k+2)(k+3) \cdots (2k-1)(2k)123 \cdots (k+1)$$

implies again $i_1 = k+1$. As in (a), $f_k(n-1; k, k+1, i_2, \dots, i_{k-1}) = f_k(n-2; k, i_2-1, \dots, i_{k-1}-1)$, and (c) follows.

For (d), consider $D_k(n-1, k-2)$ along with a similar argument to the one in Lemma 2.1(b):

$$D_k(n-1, k-2) = \sum_{k+1 \leq i_1 < \dots < i_{k-2} \leq n} f_k(n; k, i_1, \dots, i_{k-2}, 1, 2) + \sum_{k+1 \leq i_1 < \dots < i_{k-1} \leq n} f_k(n; k, i_1, \dots, i_{k-2}, 1, i_{k-1}),$$

Part (d) follows from here as in the proof of (2.4) and (2.5).

Finally, (a)–(d) yield (e). $\square$

2.4. Proof of Theorem 1.3. By (2.2), $A_k(n, k-1) - A_k(n, k) = C_k(n, k-1)$. Replacing n by $n-1$ and subtracting yields

$$\begin{aligned} (A_k(n, k-1) - A_k(n, k)) - (A_k(n-1, k-1) - A_k(n-1, k)) &= C_k(n, k-1) - C_k(n-1, k-1), \\ \Rightarrow A_k(n, k-1) - A_k(n-1, k-1) &= A_k(n, k) - A_k(n-1, k) + C_k(n, k-1) - C_k(n-1, k-1). \end{aligned}$$

By Proposition 2.6(b) and Proposition 2.7(e),

$$\begin{aligned} (2.5) \quad A_k(n, k-1) - A_k(n-1, k-1) &= A_{k-2}(n-3, k-2) + A_{k-1}(n-3, k-2) \\ &\quad + A_{k-2}(n-4, k-2) + A_{k-1}(n-4, k-2). \end{aligned}$$

The result of Corollary 2.5 completes the proof of Theorem 1.3. $\square$

2.5. Proof of Corollary 1.4. Since $f_k(n) = c_n = \frac{1}{n+1} \binom{2n}{n}$ for all $n = 0, 1, \dots, 2k-1$, we have

$$\sum_{n \geq 0} L_n^k(s) x^n = \sum_{i=0}^s (-x)^i \binom{s-i}{i} \left(f_k(x) - \sum_{j=0}^{s-i-1} x^j c_j \right).$$

Recall that the Chebyshev polynomials of the second kind satisfy the relation

$$x^{s/2} U_s \left(\frac{1}{2\sqrt{x}} \right) = \sum_{i=0}^s (-x)^i \binom{s-i}{i}$$

(see [Ri, page 75-76]), while the Catalan numbers satisfy the relation

$$\sum_{i=0}^l (-1)^i \binom{s-i}{i} c_{l-i} = (-1)^l \binom{s-1-l}{l}$$

for all $l \leq s-1$ (see [Ri, page 152-154]), and hence

$$\sum_{i=0}^s (-x)^i \binom{s-i}{i} \sum_{j=0}^{s-i-1} x^j c_j = \sum_{l=0}^{s-1} x^l \sum_{i=0}^l (-1)^i \binom{s-i}{i} c_{l-i} = \sum_{l=0}^{s-1} (-x)^l \binom{s-1-l}{l}.$$

Therefore

$$\sum_{n \geq 0} L_n^k(s) x^n = x^{s/2} U_s \left(\frac{1}{2\sqrt{x}} \right) - x^{(s-1)/2} U_{s-1} \left(\frac{1}{2\sqrt{x}} \right).$$

Finally, the Chebyshev polynomials of the second kind satisfy also the relation

$$U_m \left(\frac{1}{2\sqrt{x}} \right) = \frac{1}{\sqrt{x}} U_{m-1} \left(\frac{1}{2\sqrt{x}} \right) - U_{m-2} \left(\frac{1}{2\sqrt{x}} \right)$$

for all $m \geq 2$, hence by Theorem 1.3 we get the desired result. $\square$

3. FURTHER RESULTS

In this section we describe several directions which generalize Theorem 1.3 by use of the same arguments as in the proofs of Theorem 1.3 and Corollary 1.4.

3.1. First generalization. For any $1 \leq d \leq k-2$, let $X_{k,d}^1$ be the set of all patterns

$$(d+1, d+2, \dots, k-1, 1, 2, \dots, j, k, j+1, j+2, \dots, d)$$

for $j = 0, 1, 2, \dots, d$, plus the pattern 321. For example, $X_{3,1}^1 = \{321, 213, 231\}$. Denote the number of permutations in $S_n(321, X_{k,d}^1)$ by $x_{k,d}^1(n)$.

Theorem 3.1. *Let $k \geq 3$ and $1 \leq d \leq k-2$. For all $n \geq k$,*

$$\sum_{j=0}^{k-1} (-1)^j \binom{k-1-j}{j} x_{k,d}^1(n-j) = \binom{n-d-1}{k-d-2},$$

and for all $0 \leq n \leq k-1$, $x_{k,d}^1(n) = \frac{1}{n+1} \binom{2n}{n}$. Thus, the generating function for $\{x_{k,d}^1(n)\}_{n \geq 0}$ is

$$\sum_{n \geq 0} x_{k,d}^1(n) x^n = \frac{U_{k-2} \left(\frac{1}{2\sqrt{x}} \right) + \frac{x^{k/2}}{(1-x)^{k-d-1}}}{\sqrt{x} U_{k-1} \left(\frac{1}{2\sqrt{x}} \right)}.$$

Proof. Similarly as before, we define $\mathcal{A}_s(n, m)$ to be the size of $S_n(321, X_{k,d}^1)$ and obtain that

$$(3.1) \quad \mathcal{A}_{d+1}(n, k-d-1) = \sum_{j=0}^{k-1} (-1)^j \binom{k-1-j}{j} x_{k,d}^1(n-j)$$

On the other hand, it is easy to verify that

$$\mathcal{A}_{d+1}(n, k-d-1) = \sum_{d+1 \leq i_1 < i_2 < \dots < i_{k-d-1} \leq n} x_{k,d}^1(n; i_1, \dots, i_{k-d-1}, 1).$$

Since our permutations avoid $X_{d,k}^1$, we set $i_{k-d-1} = n$. Thus, if $\pi \in X_{k,d}^1(n; i_1, \dots, i_{k-d-2}, n, 1)$ with $d+1 \leq i_1 < \dots < i_{k-d-2} \leq n-1$, then $\pi_{k-d+1} < \dots < \pi_n$. This means

$$(3.2) \quad \mathcal{A}_{d+1}(n, k-d-1) = \binom{n-d-1}{k-d-2}$$

Combining (3.1) and (3.2) yields the desired recursive relation. The rest of the theorem is easy to derive by use of the same argument as in the proof of Corollary 1.4. $\square$

Example 3.2. For $d = 1$ and $k = 3$, Theorem 3.1 yields

$$x_{3,1}^1(n) - x_{3,1}^1(n-1) = 1, \text{ and } x_{3,1}^1(0) = x_{3,1}^1(1) = 1, x_{3,1}^1(2) = 2.$$

Hence, $x_{3,1}^1(n) = n$ for all $n \geq 1$ (see [SS]). For $d = 1$, $k = 4$ and $n \geq 0$, Theorem 3.1 yields

$$x_{4,1}^1(n) = |S_n(321, 2341, 2314)| = 2^n - n,$$

while for $d = 2$, $k = 4$ and $n \geq 2$:

$$x_{4,2}^1(n) = |S_n(321, 3412, 3142, 3124)| = 3 \cdot 2^{n-2} - 1.$$

3.2. Second generalization. Let $X_{k,d}^2$ consist of the three patterns 321, $(d+1)(d+2)\cdots(k-1)1k23\cdots d$, and $(d+1)(d+2)\cdots(k-1)k12\cdots d$. The number of $X_{k,d}^2$ -avoiding permutations in S_n is denoted by $x_{k,d}^2(n)$. Similarly as in Theorem 3.1, along with the proof arguments of Theorem 1.3 and Corollary 1.4), we arrive at

Theorem 3.3. *Let $k \geq 4$ and $2 \leq d \leq k-2$. Then for all $n \geq k$,*

$$\sum_{j=0}^{k-1} (-1)^j \binom{k-1-j}{j} x_{k,d}^2(n-j) = \sum_{j=0}^{k-1} (-1)^j \binom{k-1-j}{j} x_{k,d}^2(n-1-j) + \sum_{j=0}^{k-4} (-1)^j \binom{k-4-j}{j} x_{k,d}^2(n-3-j),$$

and $x_{k,d}^2(n) = \frac{1}{n+1} \binom{2n}{n}$ for all $0 \leq n \leq k-1$. Thus, the generating function for $\{x_{k,d}^2(n)\}_{n \geq 0}$ is

$$\sum_{n \geq 0} x_{k,d}^2(n) x^n = \frac{U_{k-1}\left(\frac{1}{2\sqrt{x}}\right) + x U_{k-3}\left(\frac{1}{2\sqrt{x}}\right)}{\sqrt{x} \left[U_k\left(\frac{1}{2\sqrt{x}}\right) + x U_{k-2}\left(\frac{1}{2\sqrt{x}}\right) \right]}.$$

Example 3.4. For $d=2$, $k=4$ and $n \geq 3$, Theorem 3.3 yields

$$x_{4,2}^2(n) = 3x_{4,2}^2(n-1) - 2x_{4,2}^2(n-2) + x_{4,2}^2(n-3), \text{ and } x_{4,2}^2(0) = x_{4,2}^2(1) = 1, \ x_{4,2}^2(2) = 2.$$

A comparison of Theorem 3.3 for different cases suggests that there should exist a bijection between the sets $S_n(X_{k,2}^2)$ and $S_n(132, X_{k,d}^2)$ for any d such that $2 \leq d \leq k-2$. Producing such a bijection remains yet an open question.

3.3. Third generalization. Let $X_{k,d}^3$ consist of the two patterns 321 and $(d+1)(d+2)\cdots(k-1)1k23\cdots d$. The number of $X_{k,d}^3$ -avoiding permutations in S_n is denoted by $x_{k,d}^3(n)$. Similarly as in the argument proofs of the main theorem in [MV3] and Theorem 1.3, we obtain

Theorem 3.5. *Let $k \geq 4$ and $2 \leq d \leq k-2$. Then for all $n \geq k$,*

$$\sum_{i=0}^k (-1)^i \binom{k-i}{i} x_{k,d}^3(n-i) = 0,$$

and $x_{k,d}^3(n) = \frac{1}{n+1} \binom{2n}{n}$ for all $0 \leq n \leq k-1$. Thus, the generating function for $\{x_{k,d}^3(n)\}_{n \geq 0}$ is

$$\sum_{n \geq 0} x_{k,d}^3(n) x^n = \frac{U_{k-1}(t)}{\sqrt{x} U_k\left(\frac{1}{2\sqrt{x}}\right)}.$$

Again, finding a direct bijection between the sets $S_n(X_{k,d}^3)$ and $S_n(132, (d+1)\cdots k12\cdots d)$ for any $k \geq 4$ and $2 \leq d \leq k-2$ is, as of now, an open question.

REFERENCES

- [Bo1] M. Bóna, The permutation classes equinumerous to the smooth class, *Electron. J. Combin.* **5** (1998) #R31.
- [Bo2] M. Bóna, The solution of a conjecture of Stanley and Wilf for all layered patterns, *J. Combin. Theory Ser. A*, **85** (1999) 96–104.
- [BW] S. C. Billey and G. S. Warrington, Kazhdan-lusztig polynomials for 321-hexagon-avoiding permutations,
- [CW] T. Chow and J. West, Forbidden subsequences and Chebyshev polynomials, *Discr. Math.* **204** (1999) 119–128.
- [Kn] D.E. Knuth, The Art of Computer Programming, 2nd ed. Addison Wesley, Reading, MA (1973).
- [Km] D. Kremer, Permutations with forbidden subsequences and a generalized Schröder number, *Discr. Math.* **218** (2000) 121–130.

- [Kr] C. Krattenthaler, Permutations with restricted patterns and Dyck paths, *Adv. in Applied Math.* **27** (2001), 510–530.
- [MV1] T. Mansour and A. Vainshtein, Restricted permutations, continued fractions, and Chebyshev polynomials *Electron. J. Combin.* **7** (2000) #R17.
- [MV2] T. Mansour and A. Vainshtein, Restricted 132-avoiding permutations, *Adv. Appl. Math.* **126** (2001), 258–269.
- [MV3] T. Mansour and A. Vainshtein, Layered restrictions and Chebychev polynomials (2000), *Annals of Combinatorics* **5** (2001), 451–458.
- [MV4] T. Mansour and A. Vainshtein, Restricted permutations and Chebyshev polynomials, *Séminaire Lotharingien de Combinatoire* **47** (2002), Article B47c.
- [R] A. Robertson, Permutations containing and avoiding 123 and 132 patterns, *Discrete Mathematics and Theoretical Computer Science*, **3** (1999) 151–154.
- [Ri] Th. Rivlin, Chebyshev polynomials. From approximation theory to algebra and number theory, John Wiley, New York (1990).
- [RWZ] A. Robertson, H. Wilf, and D. Zeilberger, Permutation patterns and continuous fractions, *Electron. J. Combin.* **6** (1999) #R38.
- [SS] R. Simion, F.W. Schmidt, Restricted Permutations, *Europ. J. of Combinatorics* **6** (1985), 383–406.
- [SW] Z. Stankova and J. West, Explicit enumeration of 321-hexagon-avoiding permutations, *Disc. Math.*, to appear.
- [W] J. West, Generating trees and forbidden subsequences, *Discr. Math.* **157** (1996), 363–372.